

Illinois Institute of Technology

PHYSICS 561
Radiation Biophysics

Andrew Howard

PHYS561 06 1/24

© 2001 Illinois Institute of Technology

Lecture 5:
Survival Models
and Curves

Andrew Howard
Illinois Institute of Technology
Physics 561: Radiation Biophysics
21 February 2001

PHYS561 06 2/24

Survival Models II

We'll present some cell-survival models that are more sophisticated than the single-hit and MTSH models; and we'll look at some results.

PHYS561 06 3/24

Overview

- **Linear-Quadratic Model**
- **Repair-Misrepair Model**
- **Lethal Potentially-lethal model**
- **Survival Curves**
- **Elkind-Sutton Experiment**
- **Repair**
- **Cell cycles**

PHYS561 06 4/24

Deficiencies in MTSH Model

- MTSH model has two problems—one at low dose and one at high:
- Doesn't predict nonzero slope near $D = 0$
- Doesn't predict nonuniform slope at high D
- Tweak for first problem:

$$S = \exp(-q_1 D) [1 - (1 - \exp(q_1 D))^n]$$
- No tweak for second problem:
 need a different model

PHYS561 06 5/24

Poisson survival

- Alpen's comment:
A cell can be killed only once, and further action on the remaining cells is constrained to that smaller number of cells
- This is equivalent to saying
 $dN = N \cdot d(f(D))$, i.e. $dN/N = d(f(D))$, or
 $\ln N = f(D) \ln N_0$ or $\ln(N/N_0) = f(D)$
- but $S = N/N_0$, so we have a basic formalism
 $\ln S = f(D)$, where $f(D)$ is a function of dose

PHYS561 06 6/24

Linear-Quadratic Model: generalized form

- Back away from mechanistic approaches and say that given Poisson statistics for lethality $\ln S = f(D)$, where f is some function
- Taylor-expand in D
 $\ln S = a_0 + a_1 D + a_2 D^2 + \dots + a_n D^n + \dots$
- We take $a_0 = 0$ because at $D = 0$ the survival fraction is 1, i.e. $\ln S = 0$.
- Thus the second-order expansion is
 $\ln S = a_1 D + a_2 D^2$

PHYS561 06 7/24

Molecular Model

- Emphasizes double-stranded breaks in DNA as the source of lasting damage
- Distinguishes between single hits causing DSBs and pairs of hits causing DSBs; ultimately the pairs of hits give rise to the quadratic dependency on D in the formulas
- Derivation in Alpen is okay, but we end up with a lot of constants that aren't independently determinable

PHYS561 06 8/24

Dual Radiation-Action Formulation

- Emphasizes that a single interaction between a high-LET radiation event and a cell produces a DSB, whereas low-LET radiation requires pairs of events
- Gives rise to a linear-quadratic model, where the one-event DSB (linear) coefficient predominates for high-LET radiation and the two-event (quadratic) coefficient predominates for low-LET radiation

PHYS561 06 9/24

Tobias's Repair-Misrepair Model


- Posits linear and quadratic mechanisms up front for repair, with explicit time-dependence
- Time-independent formulas arise at times that are long compared with cell-cycle times
- In those cases
$$S = \exp(-\alpha D)(1 + \alpha D/\epsilon)^{-1}$$
where $\epsilon = \lambda/k$ is the ratio of the repair rates for linear damage to quadratic damage.
This gives roughly quadratic behavior for $\ln S$.

PHYS561 06 10/24

Lethal—Potentially Lethal Model

- Sets up three-state system: Undamaged (A), potentially-lethally-damaged (B), and lethally damaged (C) cells
- Eupair returns state B to state A
- B automatically becomes C at long times
- Gives rise to explicit quadratic formulation
 $\ln S = \alpha D + \beta D^2$
 with α and β having explicit time-dependence

PHYS561 06 11/24



PHYS561 06 12/24

what happens if

$$D = \frac{a_1}{a_2} ?$$

$$\ln S = a_1 D + a_2 D^2$$

$$= a_1 \frac{a_1}{a_2} + a_2 \left(\frac{a_1}{a_2} \right)^2$$

$$\ln S = \frac{a_1^2}{a_2} + \frac{a_1^2}{a_2} = \frac{2a_1^2}{a_2}$$

so the quadratic and linear contributions to $\ln S$ are equal.

if $D < \frac{a_1}{a_2}$, linear mechanism predominant

but if $D > \frac{a_1}{a_2}$, quadratic mechanism predominant

PHYS561 06 13/24

Clonogenic survivability

- Even the unirradiated cells don't provide 100% survival;
- The survival for the irradiated cells has to be normalized against what's happening to the controls
- That sets an upper limit on the accuracy of the determinations
- You also need to set up a lot of plates! (Poisson statistics)

PHYS561 06 14/24

Feeder cells

- Irradiate some cells enough that they can't divide but they can metabolize
- Plate out your cells you're going to study on top of those
- This gives you a layer to feed the cells under study
- Recently, other methods have been developed


PHYS561 06 15/24

1. Errata in ch. 8:

p.169, Q2, 1st sentence

"Until the later 1950's it was not possible to use ..."

2. Fig 8.1, p173:




PHYS561 06 16/24

Shoulder of the
survival curve



PHYS561 06 17/24




Elkind-Sutton experiment:

1. Irradiate fresh cells survival curve.
2. Take cells remaining at $S \approx 0.1$ and subject them to further radiation at varying dose wounds after reaching $S = 0.1$.

PHYS561 06 18/24


RESULTS of an Ideal-Suction
experiment (fig. 8.3, p 176)



or both results of the same kind

PHYS561 06 19/24


or both results of the same kind



PHYS561 06 20/24

The Cell Cycle

Cells have a definable cycle over which specific activities occur.
Particular activities are limited to specific parts of the cycle.



Prokaryotic (PS3): characterized by (one) specific phases:

M mitosis


S synthesis (DNA & being synthesized) (replicated)

G₁ before S, after M

PHYS561 06 21/24



	how long? (hrs)	
	CHO	Hla
M	1	1
G1	1	12
S	6	8
G2	3	4
total	11	25



PHYS561 06 22/24

