

Illinois Institute of Technology

PHYSICS 561 RADIATION BIOPHYSICS

Andrew Howard

PHYS561 03 1/33

© 2001 Illinois Institute of Technology

Housekeeping

The homework associated with last week's lecture is due at 11:59 pm Friday 2 Feb. Homework for the internet students is due one week later. This pattern will persist throughout the semester. Remember that the course's internal website is

<http://icarus.csrr.iit.edu/radbio/>

and the material there is updated frequently.

Homework may be turned in either on paper, by e-mail, or by fax. Faxes for Prof. Howard:


630-252-0521

312-567-3576

Prof. Howard's pager number is 312-902-9816.

PHYS561 03 2/33

Interaction of Photons with Matter



- ◆ Let N_0 = Number of photons in
- ◆ N = Number of photons out
- ◆ t = thickness of absorber
- ◆ μ = attenuation coefficient (dimensions: L^{-1})
- ◆ Then: $N = N_0 \exp(-\mu t)$
- ◆ Mass attenuation coefficient μ/ρ (ρ = density)
- ◆ Since dimensions of density are ML^{-3} , μ/ρ has dimensions of L^2M^{-1} ; it's a *cross-section*

PHYS561 03 3/33

Cross-Section and Attenuation

Attenuations described in terms of
 $L^2/(\text{something})$

Area \rightarrow cross section

large cross section \Rightarrow high probability of
 interaction

Thus several kinds of attenuation coefficients:

Type	Dimensions	Units
◆ Linear(μ)	L^{-1}	m^{-1}
◆ Mass (μ/r)	L^2M^{-1}	m^2kg^{-1}
◆ Electronic ($_e\mu$)	L^2q^{-1}	m^2e^{-1}
◆ Atomic ($_a\mu$)	$L^2(\text{atom})^{-1}$	$m^2(\text{atom})^{-1}$

PHYS561 03 4/33

Energy Transferred and Absorbed

Energy in, out, absorbed, and leaving:

$$E_{in} \rightarrow E_{tr} + E_{out}$$

$$E_{tr} = E_{abs} + E_{leave}$$


so transferred energy is greater than absorbed energy

We define separate attenuation coefficients:

- ◆ Energy transfer attenuation coefficient
- ◆ Energy absorbed attenuation coefficient

PHYS561 03 5/33

Photoelectric Effect




Most significant at low to intermediate photon energies (~ 10 - 100 keV)

PHYS561 03 6/33


Cross-section falls off rapidly with Energy and is Z-dependent

K and L orbital edges fall within the plot for many metals, not for light atoms




PHYS561 03 7/33

Compton Scattering



1. Energy conservation
2. Momentum conservation in x
3. Momentum conservation in y



Energy conservation:

$$E_{e^-, \text{rest}} + E_{\gamma, \text{in}} = E_{\gamma, \text{out}} + E_{e^-, \text{out}}$$

$$m_0 c^2 + h\nu = h\nu' + \frac{m_0 c^2}{\sqrt{1 - v^2/c^2}}$$

Momentum conservation in x and y:

$$\frac{h\nu}{c} \hat{x} = \frac{h\nu'}{c} (\hat{x} \cos \theta + \hat{y} \sin \theta) + \frac{m_0 v}{\sqrt{1 - v^2/c^2}} [\cos \phi \hat{x} - \sin \phi \hat{y}]$$

PHYS561 03 8/33

Compton Scattering: Equations

$$\hat{x}: \quad \frac{h\nu}{c} = \frac{h\nu'}{c} \cos\theta + \frac{m_0\nu}{\sqrt{1-\nu^2/c^2}} \cos\phi$$

$$\hat{y}: \quad \frac{h\nu'}{c} \sin\theta - \frac{m_0\nu}{\sqrt{1-\nu^2/c^2}} \sin\phi$$

$$\text{Energy:} \quad h\nu - h\nu' = \frac{1}{\sqrt{1-\nu^2/c^2}} - 1$$

$$\rightarrow \alpha = \left[\frac{h\nu}{m_0c^2} \right]$$

Since $m_0c^2 = 0.511 \text{ MeV}$,
if $h\nu = 5.11 \text{ MeV}$, then $\alpha = 10$.

PHYS561 03 9/33

Compton Scattering: Results

$$KE_{out}^{(e)} = \frac{h\nu\alpha(1 - \cos\theta)}{1 + \alpha(1 - \cos\theta)}$$

$$h\nu' = h\nu - KE_{out}^{(e)}$$

$$h\nu' = h\nu - h\nu \left(\frac{\alpha(1 - \cos\theta)}{1 + \alpha(1 - \cos\theta)} \right)$$

$$= h\nu \left[\frac{1 + \alpha(1 - \cos\theta) - \alpha(1 - \cos\theta)}{1 + \alpha(1 - \cos\theta)} \right]$$

$$h\nu' = h\nu / (1 + \alpha(1 - \cos\theta))$$

PHYS561 03 10/33

Compton Scattering: Special Cases

- ◆ Recoil: $\theta = 180^\circ$, $\cos\theta = -1$
 $KE_{out}^e = 2h\nu\alpha / (1 + 2\alpha)$;
 $h\nu' = h\nu / (1 + 2\alpha)$; max. energy to e^- .
- ◆ Minimum energy transfer to electron:
 $\theta = 0^\circ$, $\cos\theta = 1$, $KE_{out}^e \approx 0$, $h\nu' = h\nu$
- ◆ Low-energy photon, $\alpha \ll 1$:
 $h\nu' \approx h\nu$, $KE_{out}^e \approx h\nu\alpha(1 - \cos\theta)$
- ◆ High-energy photon, $\alpha \gg 1$:
 $h\nu' \approx h\nu / [\alpha(1 - \cos\theta)]$, $KE_{out}^e \approx h\nu$
 (minor significance)

PHYS561 03 11/33

Compton Cross Section: Klein-Nishina formulation

diff: $\frac{d\sigma}{d\Omega} = \frac{d\sigma_o}{d\Omega} F_{KN}$ where $\frac{d\sigma_o}{d\Omega} = \frac{r_o^2}{2} (1 + \cos^2 \theta)$
Thompson differential x - section

\uparrow
 Thompson scatt cross section

\downarrow
 total cross $\Rightarrow \int \frac{d\sigma}{d\Omega} d\Omega$

$r_o =$ classical electron radius = $2.8 \cdot 10^{-15}$ M
 $= \frac{ke^2}{m_o c^2}$ where k is coulomb-law constant

total cross-section for Thompson case:

$$\begin{aligned} \sigma_{tot} &= \int_{\text{all solid angle}} \frac{d\sigma_o}{d\Omega} d\Omega = \frac{r_o^2}{2} \int_0^\pi \int_0^{2\pi} (1 + \cos^2 \theta) \sin\theta d\phi d\theta \\ &= \frac{r_o^2}{2} 2\pi \int_0^\pi (1 + \cos^2 \theta) \sin\theta d\theta \\ &= \pi r_o^2 \int_0^\pi (1 + \cos^2 \theta) \sin\theta d\theta \Rightarrow \begin{matrix} u = \cos\theta \\ du = -\sin\theta d\theta \end{matrix} \end{aligned}$$

PHYS561 03 12/33

Formulae for Cross-Sections

Total Cross-section:

$$\begin{aligned}\tau_{\text{tot}} \sigma_o &= \pi r_o^2 \int_1^{-1} (1 + u^2) (-du) \\ &= -\pi r_o^2 \left(u + \frac{u^3}{3} \right) \Big|_1^{-1} \\ &= -\pi r_o^2 \left[-1 - 1 + \left[\left(-\frac{1}{3} \right) - \left(\frac{1}{3} \right) \right] \right] \\ \tau_{\text{tot}} \sigma_o &= -\pi r_o^2 \left[-2 - \frac{2}{3} \right] = \pi r_o^2 \left(2 + \frac{2}{3} \right) = \frac{8\pi r_o^2}{3} \\ \tau_{\text{tot}} \sigma_o &= \frac{2}{3} (SA)_{r_o}, \text{ where } S.A. = \text{surface area} \\ &\quad \text{of sphere } 4\pi r_o^2\end{aligned}$$

PHYS561 03 13/33

Compton Scattering Cross Section

$$4.27: \frac{d_e \sigma_I}{d\Omega} = \frac{r_o^2}{2} (1 + \cos^2 \theta)$$

$$4.29: \frac{d\sigma}{d\Omega} = \frac{d\sigma_o}{d\Omega} F_{KN} = \frac{r_o^2}{2} (1 + \cos^2 \theta) F_{KN}$$

F_{KN} is a geometry-dependent quantum-mechanical factor given as eqn. 4.30;


$F_{KN} \leq 1$; $F_{KN} \rightarrow 1$ as $\theta \rightarrow 0^\circ$ or as $\alpha \rightarrow 0$.

PHYS561 03 14/33

Energy of Compton Electrons

- ◆ From standard Klein-Nishina equations we can determine the spectrum of Compton electron energies.
- ◆ KE_{\max} of electron is close to photon energy
 - $E_{\gamma} = 0.5 \text{ MeV}$ implies $KE_{\max}(e^{-}) = 0.331 \text{ MeV}$
 - $E_{\gamma} = 1.0 \text{ MeV}$ implies $KE_{\max}(e^{-}) = 0.796 \text{ MeV}$

PHYS561 03 15/33



PHYS561 03 16/33

Effect of Binding Energy


- ◆ Typical Compton treatments assume free electrons: this is close to right.
- ◆ Sharp fall-off in total coefficient at low energies (below 50 KeV), but not much gets transferred at those energies anyway, it doesn't affect the equations much

PHYS561 03 17/33

Pair Production

- ◆ Can happen if $E_\gamma > 1.022 \text{ MeV} = 2 * m_0(e^-)$
- ◆ Rapidly increasing cross-section $> 1.022 \text{ MeV}$
- ◆ Stopping power/atom varies as Z^2
- ◆ Energy transferred is $(h\nu - 1.022) \text{ MeV}$
- ◆ Scattering nucleus plays fairly passive role (not much momentum transferred to nucleus)
- ◆ Generally the positron gets annihilated, giving off another pair of 0.511 MeV photons. These generally escape and are not part of the absorbed energy

PHYS561 03 18/33



PHYS561 03 19/33

Bremsstrahlung: Radiative Energy Loss

- ◆ “Braking radiation”:
A fast electron loses energy to its environment in a nonspecific way due to Coulombic interaction with neighboring charged particles.
- ◆ The static particles are much more massive than the electron, so they don’t get accelerated nearly as much as the electron does: but the electron does get accelerated.
- ◆ What happens when an electron is accelerated? It has to radiate! This type of Coulombically-motivated radiation is *Bremsstrahlung*


PHYS561 03 20/33

Significance of Bremsstrahlung

- ◆ Example in X-ray generators:
- ◆ 1.5418\AA (8KeV) X-rays are produced in great quantity when we shoot fast electrons at a copper target
- ◆ BUT: we also get a lot of radiative transfer of energy from the electrons as they move past the copper atoms. This gives rise to Bremsstrahlung, which has no characteristic energies.
- ◆ Thus the spectrum is like this:

PHYS561 03 21/33

Output X-ray Spectrum of a Copper Target



PHYS561 03 22/33

Energy Transfer

- ◆ Compton Processes in Tissue
- ◆ Charged Particles and Matter
- ◆ Final Steps in Energy Absorption
- ◆ Dose and Kerma: A Review
- ◆ Neutron Interactions with matter

PHYS561 03 23/33

Compton Processes in Tissue


Biological soft tissue is predominantly made up of H, C, N, O, and a little P and S. So attenuation of photons is dominated by those light elements ($Z \leq 16$)

PHYS561 03 24/33

Interaction of Charged Particles with Matter

See pages 84 & 85 in the text-

Provides solutions to the dynamical equations describing motion of a heavy charged particle past a stationary electron or (by relativity) motion of an electron past a stationary heavy particle: $F = kze^2/r^2$ along line MQ



PHYS561 03 25/33

Interaction of e^- With Heavy Charged Particle

$$r_o = \frac{ke^2}{m_o c^2}$$

$$F = \frac{kq_M q_e}{r^2} = \frac{kze \cdot e}{r^2}$$

- ◆ Momentum imparted to electron associated with force:

$$\vec{p} = \int_{-\infty}^{\infty} \vec{F} dt \quad \text{classical coherent}$$

$$p_y = \int_{-\infty}^{\infty} F_y dt = kze^2 \int_{-\infty}^{\infty} \cos \theta dt = \frac{2zr_o m_o c^2}{vb}$$

non relativistic


$$\Delta E = \frac{(\Delta p)^2}{2m_o} = \frac{z^2 r_o^2 m_o c^4 M}{b^2 E} \quad \left(E = \frac{1}{2} Mv^2 \right)$$

PHYS561 03 26/33

Dose and Kerma

See Fig. 5.5 in text.

Because secondary events extend farther into tissue (or other) than the initial deposited radiation, dose extends farther into the interior than kerma.




PHYS561 03 27/33

Neutrons: Elastic Scatter

Important up to ~14 MeV range

Energy imparted to nucleus:



$$E_t = E_n \frac{4M_n M_A}{(M_A + M_n)^2} \cos^2 \theta$$

average over angles:

$$\text{average} [f(x)]_a^b \equiv \frac{1}{b-a} \int_a^b f(x) dx$$

$$\bar{E}_T = \frac{E_n \cdot 2M_n M_A}{(M_n + M_A)^2}$$

PHYS561 03 28/33

Average of function: $\langle f(x) \rangle_a^b = \frac{1}{b-a} \int_a^b f(x) dx$

$$f(x) = \cos^2 x, \quad a=0, \quad b=2\pi$$

$$\langle \cos^2 \theta \rangle = \frac{1}{2\pi - 0} \int_0^{2\pi} \cos^2 \theta d\theta = \frac{1}{2\pi} \int_0^{2\pi} \cos^2 \theta d\theta.$$

Now $\cos^2 \theta - \sin^2 \theta = \cos 2\theta \leftarrow$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

In this case let $A=B=\theta$:

$$\cos(\theta+\theta) = \cos 2\theta = \cos \theta \cos \theta - \sin \theta \sin \theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

but $\cos^2 \theta + \sin^2 \theta = 1$; $\sin^2 \theta = 1 - \cos^2 \theta$

$$\cos 2\theta = \cos^2 \theta - (1 - \cos^2 \theta)$$

$$\cos 2\theta = 2\cos^2 \theta - 1$$

PHYS561 03 29/33

$$\cos 2\theta = 2\cos^2 \theta - 1$$

$$\frac{\cos 2\theta + 1}{2} = \cos^2 \theta$$

$$\therefore \langle \cos^2 \theta \rangle = \frac{1}{2\pi} \int_0^{2\pi} \cos^2 \theta d\theta$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \frac{\cos 2\theta + 1}{2} d\theta$$

$$= \frac{1}{2\pi} \left[\int_0^{2\pi} \frac{\cos 2\theta}{2} d\theta + \int_0^{2\pi} \frac{1}{2} d\theta \right]$$

$\theta = \frac{1}{2}u$
let $u = 2\theta$, $du = 2d\theta$
 $\frac{1}{2}du = d\theta$

$$= \frac{1}{2\pi} \left[\int_0^{2\pi} \frac{\cos u}{2} \frac{du}{2} + \frac{1}{2} \int_0^{2\pi} d\theta \right]$$

PHYS561 03 30/33

$$= \frac{1}{2\pi} \left[\frac{1}{4} \int_0^{2\pi} \cos u \, du + \frac{1}{2} \cdot 2\pi \right]$$

$$= \frac{1}{2\pi} \left[\frac{1}{4} \sin u \Big|_0^{2\pi} + \pi \right]$$

$$\langle \cos^2 \theta \rangle = \frac{1}{2\pi} [0 + \pi] = \frac{1}{2}$$

PHYS561 03 31/33



Inelastic Scatter

Increasingly important at higher neutron energies



PHYS561 03 32/33

Neutrons: Other Mechanisms

(III) Nonelastic (75 MeV)



(IV) Neutron Capture



(V) Spallation: Nucleus fragments!

Need very high-energy neutrons ($> 100 \text{ MeV}$)