

# **Illinois Institute of Technology**

---

## Physics 561 Radiation Biophysics

Andrew Howard

PHYS561 14 1/28

© 2001 Illinois Institute of Technology

## **Class Overview**

---

- ◆ Genetics II
  - Review of definitions
  - Frameshifts and substitutions
  - Relative sensitivities of cell types
  - Organismal differences
  
- ◆ High-LET Radiation
  - Review of definitions
  - Physics of energy deposition
  - Impact on DNA
  - Resistance to repair

PHYS561 14 2/28

## Frameshifts

---

- Most common form of DNA damage
- Chemistry
  - Deletion of bases
  - Fragmentation of sugar-phosphate backbone
  - Distortion of base-base hydrogen bonds and other 3-D elements
- Results in complete misreading of remainder of message

PHYS561 14 3/28

## Genes

---

- ◆ Unit of DNA that codes for a specific function
- ◆ Contain exons and introns
- ◆ Genes and the Processes for which DNA is Responsible

PHYS561 14 4/28

## Structural elements

---

- ◆ Bases
  - Adenine
  - Cytosine
  - Guanine
  - Thymine
- ◆ Phosphate-deoxyribose backbone

PHYS561 14 5/28

## Special Features

---

- ◆ Code is redundant
- ◆  $4^3 = 64$  codons
- ◆ 20 amino acids + 1 or 2 control codes; so most amino acids have more than one codon associated with them.
- ◆ Middle base is conserved, i.e. all the codons for a given amino acid have the same middle base.
- ◆ First RNA base can be sloppy

PHYS561 14 6/28

## Errors in Fidelity and Their Consequences

---

- ◆ How do Chemicals & Radiation Affect Fidelity (rate of mutation)?
- ◆ Increase likelihood of replication error
- ◆ Radiation: bond disruption in bases (or sugars)
  - Direct
  - Indirect

PHYS561 14 7/28

## Repair

---

- ◆ Cell has mechanisms to recognize & replace faulty bases before they have a chance to be replicated
- ◆ Some injury may disrupt a large enough segment of DNA that repair either fails or is error-prone

PHYS561 14 8/28

## High LET Radiation

---

- ◆ Review definition of LET:  
Rate of change of energy with distance along a track
- ◆ Caveat I: Local Deposition
- ◆ Caveat II: LET only assoc. with charged particles

PHYS561 14 9/28

## LET Equation: Bethe-Block Formulation

---

- ◆ If we let:
  - $z^*$  = Effective charge of projectile particle
  - $Z$  = Atomic number of atoms in medium
  - $A$  = Atomic weight of atoms in medium
  - $C$  = sum of electron shell corrections
  - $\delta/2$  = condensed medium correction
  - $\beta = v/c$  for particle
- ◆  $-dE/dx = (0.307z^{*2}Z/A\beta^2) \cdot (\ln(2m_0c^2\beta^2)/(1-\beta^2)) - \beta^2 - C/Z - \delta/2$

PHYS561 14 10/28

## Where is Energy Deposited?

---


- ◆ Most of the energy is deposited just before the particle stops moving
- ◆ We're interested in *average* LET over a region or track

PHYS561 14 11/28



## Calculating Averages of Continuous Variables

---

We wish to calculate the mean of a continuous function  $f(x)$  over a range of  $x$  from  $a$  to  $b$ ,  
Then  $\langle f(x) \rangle_{a,b} = [\int_a^b f(x) dx] / (b-a)$



PHYS561 14 12/28



## Average LET, keV $\mu$ m<sup>-1</sup>

---

Table 14.1:

<i>Radiation type</i>	$(LET_{av})_T$	$(LET_{av})_E$
<sup>60</sup> Co $\gamma$ -rays	0.27	19.6
250 kVp x-rays	2.6	25.8
3 MeV neutrons	31	44
Radon $\alpha$ rays	118	83
14 MeV neutrons	11.8	125
Recoil protons	8.5	25
Heavy recoils	142	362

PHYS561 14 15/28

## Direct & Indirect Effects

---

◆ High LET means that many ionizations occur in a small neighborhood

	LET Kev/ $\mu$ m	Spur energy eV	Events/ $\mu$ m	Spacing nm
<sup>60</sup> Co $\gamma$	0.25	60	4	250
Radon $\alpha$	118	60	2000	0.5

◆ This fact by itself accounts for much of the difference in biological consequence of high LET radiation

PHYS561 14 16/28

## Biological Effects of High-LET Radiation

---


- ◆ RBE = relative biological effectiveness
- ◆ analogous to OER in its definitional form:  
$$RBE = \frac{\text{dose for given end point for reference radiation}}{\text{dose for given end point for test radiation}}$$
- ◆ Problem with this formulation:
  - assumes that dose-response curves are described by identical functions, i.e.
  - RBE is independent of the response level at which it is estimated. Often not true!

PHYS561 14 17/28

## Relationship between RBE and LET

---

- ◆ RBE vs LET:
- ◆ Generally higher LET radiations have higher RBE, up to a certain point.
- ◆ With some systems the curve turns over.



PHYS561 14 18/28


## Cell-Cycle Dependence for RBE

- ◆ Low LET radiations exert much more effect in late G2 & M than elsewhere
- ◆ High LET radiation exerts approximately equal effects throughout the cell cycle
- ◆ Reason: irreparable damage early in the cycle will persist through mitosis, so it's just as bad!

PHYS561 14 19/28

## OER vs LET

- ◆ OER for High-LET Radiation is generally smaller than for low-LET radiation because the damage is less dependent on oxygen fixation of radical species.
- ◆ This plot shows OER going not to 1 but to ~1.5--because water-derived radicals are still produced at high LET; cf. Fig. 14.6.



PHYS561 14 20/28

## LET vs Fractionation

---

- ◆ Recall that repair-competent systems respond less to fractionated doses than to single doses whereas repair-deficient systems are fractionation-independent
- ◆ For high-LET radiation fractionation does not decrease biological effects
- ◆ Sometimes in tumors fractionation increases biological effect!
  - Why? - no explanation in Alpen
  - 2-stage model for cancer: (Ullrich)
  - Also: high dose rate is more likely to simply kill the cell rather than producing clonogenically competent but mutated cells

PHYS561 14 21/28

## Late Effects of High LET Radiation


---

- ◆ You can induce cancer with neutrons
- ◆ Ignores fractionation or (!) is more likely with fractionation
- ◆ Tumors also arise from heavy-ion irradiation


PHYS561 14 22/28

## LET vs. Cross Section


- ◆ Alpen looked at particle fluence as a dose parameter--requires constant LET over volume
- ◆ Result: power-law relationship between cross section for tumor induction and LET (fig.14.7)



General Note  
re linearity



PHYS561 14 24/28



$a \ln x$   
 $y = ke^{a \ln x}$   
 but  $a \ln x = \ln x^a$   
 $y = ke^{\ln x^a} = kx^a$   
 $y = kx^a$  power law

PHYS561 14 26/28

## Probability of Traversals

◆ Table 14.2 shows that multiple traversals of the nucleus are very rare even at high doses

◆ It also shows a close correspondence between cross section for tumor production and cross-section for traversal

Dose, Gy	<fluence/ cell>	Probability of traversals			
		0	1	2	>= 1
0.01	0.006	0.993	0.006	$2.1 \times 10^{-5}$	0.006
0.02	0.013	0.987	0.013	$6.8 \times 10^{-5}$	0.013
0.05	0.032	0.968	0.031	$5.0 \times 10^{-4}$	0.032
0.15	0.097	0.907	0.088	$4.0 \times 10^{-3}$	0.092
0.20	0.129	0.878	0.114	$7.0 \times 10^{-3}$	0.121
0.30	0.193	0.823	0.159	$1.5 \times 10^{-2}$	0.176
0.40	0.258	0.772	0.199	$2.5 \times 10^{-2}$	0.227

PHYS561 14 27/28

## Cell Transformation in High-LET Irradiation

- ◆ Assertion: non local effects on DNA predominate over point mutation events
- ◆ Certain assay systems suggest this assertion:
- ◆ Kronenberg et al, human TK6 lymphoblasts--entire *hprt* gene is missing after irradiation
- ◆ "If large genomic changes are brought about by heavy ion radiation, then a multistep process may be short circuited to a single event."

PHYS561 14 28/28