

Illinois Institute of Technology

Physics 561
Radiation Biophysics

Andrew Howard

PHYS561 07 1/31

© 2001 Illinois Institute of Technology

Illinois Institute of Technology

Physics 561
Radiation Biophysics:
Modifiers of the Radiation Response
Andrew Howard
BCPS Department
28 February 2001

PHYS561 07 2/31

Class Plan

- ◆ Review Midterm
- ◆ Biological Modifiers of Radiation Response
- ◆ Break
- ◆ Chemical/Physical Modifiers of Radiation Response

PHYS561 07 3/31

Midterm, Problem 1

Half-life of ^{32}P = 14.29 days

$$\tau = \text{Mean life} = T_{1/2} / 0.693 = 14.29 / 0.693 \text{ days} \\ = 20.62 \text{ days}$$

$$\text{Decay constant} = 0.693 / T_{1/2} = 0.0485 \text{ day}^{-1} \\ = 0.0485 \text{ day}^{-1} / 86400 \text{ sec/day} = 5.613 \cdot 10^{-7} \text{ sec}^{-1}$$

after 10 days: $A = A_0 \exp(-\lambda t)$

$$A = 3.000 \text{ mCi} \cdot \exp(-0.0485 \text{ day}^{-1} \cdot 10 \text{ day}) = \\ 3.000 \cdot \exp(-0.485) = 1.847 \text{ mCi} = \\ 1.847 \cdot 3.700 \cdot 10^7 \text{ Bcq} = 6.834 \cdot 10^7 \text{ Bcq.}$$

PHYS561 07 4/31

Midterm, Problem 2

Eqn. 4.20a and $v = c/\lambda$ means that

$$hc/\lambda' = [hc/\lambda] \gamma (1 + \alpha(1 - \cos\theta))$$

$$\text{so } \lambda'/\lambda = 1 + \alpha(1 - \cos\theta)$$

$$\text{Thus } (\lambda'/\lambda - 1)/\alpha = 1 - \cos\theta$$

$$\cos\theta = 1 - (\lambda'/\lambda - 1)/\alpha$$

But $\lambda' > \lambda$: the photon has to lose energy.

$$\lambda' = \lambda * 1.0001; \lambda'/\lambda = 1.0001, \lambda'/\lambda - 1 = 10^{-4}$$

$$\text{now } \alpha = h\nu/m_0c^2 = hc/(\lambda m_0c^2).$$

$$\text{For } \lambda \text{ in nm, } E \text{ in eV, } E = hc/\lambda = 1239.84 / \lambda$$

$$hc/\lambda = 1239.84 / 0.15 \text{ eV} = 8265.6 \text{ eV} =$$

$$= 8.2656 * 10^{-3} \text{ MeV}$$

$$\alpha = hc/(\lambda m_0c^2) = 8.2656 * 10^{-3} / 0.511 = 0.01618$$

PHYS561 07 5/31

Midterm problem 2, concluded

$$\cos\theta = 1 - (\lambda'/\lambda - 1)/\alpha = 1 - 10^{-4} / 0.01618$$

$$\text{so } \cos\theta = 1 - 0.006180 = 0.99382$$

$$\text{We can compute } \theta = \arccos(0.99382) = 6.373^\circ.$$

We can also compute it by an expansion:

$$\text{To first order } \cos(x) = 1 - x^2/2 \text{ so}$$

$$1 - \theta^2/2 = 0.99382,$$

$$\text{i.e. } \theta^2/2 = 0.006180, \theta^2 = 0.0123609,$$


$$\theta = 0.11118 \text{ rad} = 6.37^\circ.$$

$$KE_{\text{max}} = \text{the missing energy} = hc/\lambda' - hc/\lambda =$$


$$(hc/\lambda) \chi 10^{-4} = 8265.6 \text{ eV} * 10^{-4} = 0.82656 \text{ eV}.$$

PHYS561 07 6/31

Midterm I, problem 3.



Eqn. 6-14, p.117



PHYS561 07 7/31


Midterm 1, problem 3, concluded

Cysteine and glutathione contain thiol (-SH) groups, which are good free-radical scavengers. They are likely to reduce the effects of radiation on cells by reacting with free radicals and producing relatively innocuous products--specifically, disulfide compounds, in which two molecules of a sulfur-containing free radical react with one another to pair up. This would then reduce the chemical damage in biological tissue that otherwise would arise from the reactions of water with the free radicals.

PHYS561 07 8/31

Midterm problem 4

Does this curve display MTSH survival characteristics? Yes, since it's more or less linear at intermediate doses and flattens out at low dose.



PHYS561 07 9/31

Midterm problem 4, continued

- ◆ Estimate n , D_0 , D_q :
 - n is the intercept on the S axis. If we extrapolate up from the linear portion we find that it hits the y axis at about $\ln(S) = 1.6$. This corresponds to $n = e^{1.6} = 4.95$.
- ◆ D_q is the dose in Grays at which the linear portion of the curve hits $S=1$, i.e. $\ln S = 0$. On the graph that appears to happen at about $D = 1.7$, so $D_q = 1.7$
- ◆ Two ways to calculate D_0 :
 - $D_0 = -1/\text{slope} = 1/[-(2.45-1.45)/(4-3)] = 1 \text{ Gy}$.
 - $D_0 = D_q/\ln n = 1.7/1.6 \sim 1 \text{ Gy}$.

PHYS561 07 10/31

Midterm, Problems 5-7

- ◆ **5. Recombination:**
Recently produced free-radical pair joins up again to reform the original molecule
 $A-B + h\nu \rightarrow A^{\cdot} + \cdot B \rightarrow A-B$ fast ($\sim 10^{-11}$ s)
- ◆ **6. 1 Gy is a unit of absorbed dose:**
1 Gy unit of energy transferred per unit mass.
1 Gy = 1 Joule of transferred energy per kilogram of mass
- ◆ **7. Hydroxyl radical**
•OH or O-H•
(or draw out the dots for all valence electrons)

PHYS561 07 11/31

Midterm, Problem 8

- ◆ **High - LET Radiation**
LET = Linear energy transfer so high - LET radiation is radiation that transfers large amounts of energy per unit linear travel or energy per thickness in a medium.
Neutrons and alpha particles are examples of high - LET radiation

PHYS561 07 12/31

Midterm Problem 9: Extra Credit

- ◆ *Would recombinational repair be as effective in dealing with lesions in the Y chromosome as it is in dealing with lesions in chromosome 4?*
- ◆ **No.** Recombinational repair requires the presence of homologous duplex DNA. There are two copies of chromosomes 1-22 in every human cell that has a nucleus, so the homologous duplex DNA is present in the "other" copy of the chromosome. The Y chromosome is present in only one copy in the male (and absent altogether in the female), so there is no homologous duplex DNA from which to make a correction.

PHYS561 07 13/31

Effectors of Radiation Sensitivity

- ◆ Biological
 - Cells go through life cycles & are much more sensitive to radiation damage at some stages than at others
- ◆ Chemical
- ◆ Physical


PHYS561 07 14/31

Cellular Life Cycles

	<u>Phases</u>	
♦ Mitotic	M	(short)
- Sensitive		
♦ Presynthetic	G ₁	(variable)
- Radiation causes 1/2 h delay here		
♦ Synthetic	S	(4 - 8 h)
- DNA synthesis		
- Least sensitive		
♦ Postsynthetic	G ₂	(usually short 1 - 2h)
- Radiation causes 3 - 4 h delay here		
- End of G ₂ sensitive		


14 h

PHYS561 07 15/31



PHYS561 07 16/31

Physical & Chemical Effectors



Pure water

$$\frac{1\text{kg}}{1\text{L}} \quad \text{MW}(\text{H}_2\text{O}) = 18$$

water 18g/mole

$$1 \text{ kg H}_2\text{O} = \frac{1000\text{g}}{18\text{g/mole}} = 55.5 \text{ moles}$$
$$\rho = \frac{1\text{kg}}{\text{L}} = 55.5 \frac{\text{moles}}{\text{L}} = 55.5 \text{M}$$

PHYS561 07 17/31

Assignment for 9 March

1. Problem 1, ch.9, Alpen

2. A human has a fever such that her body temperature is 39°C. This fever is accompanied by an increase in white-blood cells. How will these conditions affect her sensitivity to radiation?

PHYS561 07 18/31


Radiation & Temperature

Kinetics


37°C = 310K
27°C = 300K

p199

T < 100K
100 < T < 170 K
170 < T < 420 K



PHYS561 07 19/31



Radiation and Temperature

◆ Effects in various temperature ranges:

- T_{min} T_{max} Effects
- 0K 100K temp-insensitive; no charge migration, target must be hit
- 100K 170K excitation localized at site; exciton migration crucial
- 170K 420K disruption of disulfides, ionization

◆ Does this matter much with biological systems, where T ~ 310K almost all the time? Probably not, other than for understanding mechanisms.


PHYS561 07 20/31

Oxygen and Radiation



- ◆ FACT: O_2 is a radiation-sensitive molecule
- ◆ H_2O free-radical chemistry in the presence of O_2 is different from H_2O free radical chemistry in the absence of O_2 .
- ◆ $P(O_2)$ in tissue varies widely
 - Hemoglobin transports O_2
 - Myoglobin stores O_2

PHYS561 07 21/31

Damage Fixation by Oxygen




Mitigators of $\left\{ \begin{array}{l} O_2 \text{ fixation} \\ \text{radical presence} \end{array} \right.$



PHYS561 07 22/31

Experiments on Oxygen Sensitivity




PHYS561 07 23/31

Oxygen Enhancement Ratio

$$\text{OER} \equiv \frac{\text{dose in } N_2 \text{ for surviving fraction } S/S_0}{\text{dose in } O_2 \text{ for surviving fraction } S/S_0}$$

if $S = S_0/10$



$$\text{OER} = \frac{D_N}{D_O}$$

PHYS561 07 24/31

Quantitative Oxygen Sensitivity

$$\frac{S}{S_N} = \frac{m[O_2] + k}{[O_2] + k}$$

k is in concentration units



m is unitless

[O₂] if [O₂] >> k, S/S_N = m


Organism	m	K, μM
Shigella	2.9	4.0
E.coli	3.1	4.7
S.cerevisiae	2.4	5.8

PHYS561 07 25/31

Thiol Mitigators




PHYS561 07 26/31



Sensitization by Nitroaromatics

How do we make the cells in the rapidly growing tumor as rad-sensitive as they would be if P_{O_2} were higher?

Nitroaromatics react with radicals to "fix" (stabilize) the damage



PHYS561 07 28/31

$$\frac{S_{21}}{S_{21}} = 2.05 \text{ at } 0.355\%$$

translate that to molarity:

$$\text{mass fraction} = 0.355\% = 3.55 \cdot 10^{-3}$$

$$\frac{\text{mass O}_2}{\text{total mass}} = 3.55 \cdot 10^{-3}$$

$$\text{mole fraction} = \frac{3.55 \cdot 10^{-3} \text{ g/gas}}{\frac{16 \text{ g/mole}}{22.4 \text{ L}}} = \frac{3.55 \cdot 10^{-3}}{0.714} = 3.1 \cdot 10^{-3}$$

$$\text{mole frac} = 3.1 \cdot 10^{-3}$$

Vol at STP 1 mole of gas $\rightarrow 22.4 \text{ L}$
 \downarrow
 O_2 soln

$$\frac{\text{mole fraction}}{22.4 \text{ L}} = \frac{3.1 \cdot 10^{-3}}{22.4} \text{ M}$$

PHYS561 07 29/31

$$\text{mole frac} = 3.1 \cdot 10^{-3}$$

Vol at STP 1 mole of gas $\rightarrow 22.4 \text{ L}$
 \downarrow
 O_2 soln

$$\frac{\text{mole fraction}}{22.4 \text{ L}} = \frac{3.1 \cdot 10^{-3}}{22.4} \text{ M}$$

$$= 0.138 \cdot 10^{-3} \text{ M} = 138 \mu\text{M}$$

PHYS561 07 30/31

$$0.355\% = 3.55 \cdot 10^{-3} \text{ g/g H}_2\text{O}$$

$$\frac{1 \text{ g}}{1000 \text{ mL}} = 10^{-3} \text{ L}$$

$$\rightarrow \frac{3.55 \cdot 10^{-3} \text{ g}}{(10^{-3} \text{ L})(1000 \text{ g/L})}$$

— Interval —

PHYS561 07 31/31